



## PATENT ABSTRACTS OF JAPAN

(11) Publication number: 11171848 A

(43) Date of publication of application: 29.06.1999

(51) Int. Cl. C07C237/40

A61K 31/00, A61K 31/00, A61K 31/16, A61K 31/19, A61K 31/38,  
 A61K 31/44, A61K 31/445, C07C255/58, C07C311/46, C07C311/51,  
 C07D213/56, C07D213/80, C07D295/14, C07D333/24, C07D333/38

(21) Application number: 10270721

(22) Date of filing: 25.09.1998

(30) Priority: 26.09.1997 JP 09277942

(71) Applicant: FUJIREBIO INC

(72) Inventor: IKAWA HIROSHI  
 NISHIMURA MASAHITO  
 OKADA KEIJI  
 NAKAMURA TAKASHI

(54) AROMATIC AMIDE DERIVATIVE

and a base in an inert solvent.

COPYRIGHT: (C)1999,JPO

(57) Abstract:

PROBLEM TO BE SOLVED: To obtain the subject new compound having Acetyl-CoA Carboxylase (hereafter, referred to as ACC) inhibitory activity, and useful for the treatment of visceral adiposis syndrome as a risk factor of geriatric diseases such as myocardial infarction, cerebral infarction and diabetes.

SOLUTION: This new compound is an aromatic amide derivative of formula I ( $R^1$  and  $R^2$  are each H, a 1-12C alkyl, aromatic hydrocarbon group, aromatic heterocyclic group or the like;  $R^3$  is H, a substituted amino, 1-12C alkyl, 2-12C alkenyl or the like; Y is  $CH=CH$ ,  $N=CH$ , or the like;  $R^4$  is an acidic functional group; ring A is an aromatic hydrocarbon group, aromatic heterocyclic group or cyclic alkyl), e.g. 2-[2-(3-trifluoromethylphenylamino)benzamido]benzoic acid. The amide derivative of formula I is obtained, for example, by condensation reaction between an amino compound of formula II and a carboxylic acid compound of formula III in the presence of a condensation agent

